



ORAL PRESENTATION

Open Access

Current Munich status concerning in-vivo optical coherence tomography for differentiating lesions of the upper aerodigestive tract

Veronika Volgger^{1*}, Herbert Stepp¹, Waseem Jerjes², Tahwinder Upile², Andreas Leunig¹, Colin Hopper², Christian Betz¹

From 2nd Scientific Meeting of the Head and Neck Optical Diagnostics Society
San Francisco, CA, USA. 23-24 January 2010

Objective

Non-invasive differentiation of pre- and early malignant mucosal changes may be helpful to reduce the morbidity and shorten the time to diagnosis for the patients concerned. Optical Coherence Tomography (OCT) seems to be well suited to reach this goal.

Methods

61 patients with a total of 82 primary leukoplakic or erythroplakic mucosal lesions of the upper aerodigestive tract (OADT) were prospectively examined using an in-vivo, time-domain OCT (Niris®, Imalux Corporation, USA; lateral resolution 25 µm / axial resolution 15 µm) so far. The results were subsequently correlated to histopathology of tissue biopsies taken from the same areas. Additionally, intensity profiles of OCT images were evaluated concerning their ability to differentiate dysplasia from hyperplasia.

Results

Down to a depth of 1.5 mm, microanatomical structures were clearly identifiable on the OCT images. Concerning the ability to differentiate non- and early invasive lesions, OCT reached a sensitivity of 100% and a specificity of 92% or 75% (investigator unblinded or blinded to visual inspection), respectively. False positive findings were mostly hyperplastic lesions with a significant broadening of the epithelial layer. Epithelial thicknesses measured on OCT images and histological slides correlated well ($\kappa=0.63$), but were no indicators of either epithelial hyperplasia or different grades of dysplasia.

Yet, OCT intensity profiles showed a statistically significant difference between dysplastic and hyperplastic lesions in a subset of cases (n=44).

Conclusions

The method seems highly promising for early, non-invasive tumour diagnosis in the UADT. Technical advances and an increase in patient numbers will help to define its clinical value in the near future.

Author details

¹Department of Otorhinolaryngology, Head and Neck Surgery, Ludwig Maximilian University, Munich, Germany. ²UCL Department of Surgery, University College London Medical School, London, UK.

Published: 29 October 2010

doi:10.1186/1758-3284-2-S1-O39

Cite this article as: Volgger et al.: Current Munich status concerning in-vivo optical coherence tomography for differentiating lesions of the upper aerodigestive tract. *Head & Neck Oncology* 2010 **2**(Suppl 1):O39.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Department of Otorhinolaryngology, Head and Neck Surgery, Ludwig Maximilian University, Munich, Germany
Full list of author information is available at the end of the article